

Generalization in the Hopfield Model

Leonid B. Litinskii

High Pressure Physics Institute
of Russian Academy of Sciences

Russia, 142092 Troitsk Moscow region,
e-mail: u10740@diapod.podolsk.ru

In the Hopfield model the ability of the network to generalization is studied in the case of the network trained by one input image (*the standard*).

Basic Model

The maximization problem with the symmetrical connection matrix is considered:

$$\begin{cases} F(\vec{\sigma}) = \sum^n J_{ij} \sigma_i \sigma_j \rightarrow \max, & \sigma_i = \{\pm 1\} \quad \forall i \\ J_{ij} = J_{ji}, & J_{ii} \text{ does not matter}^1. \end{cases} \quad (1)$$

The configuration vector $\vec{\sigma} = (\sigma_1, \sigma_2, \dots, \sigma_n)$ providing the solution of the problem is called the ground state.

The Hebbian-like representation exists for every $\mathbf{J}$,

$$\mathbf{J} = \mathbf{S}^T \cdot \mathbf{S}, \quad \text{where } \mathbf{S} \text{ is a real } (p \times n)\text{-matrix and } p = \text{rank } \mathbf{J}. \quad (2)$$

We would like to investigate the special case of the $(p \times n)$ -matrix $\mathbf{S}$:

$$\mathbf{S} = \begin{pmatrix} 1-x & 1 & \dots & 1 & 1 & \dots & 1 \\ 1 & 1-x & \dots & 1 & 1 & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \dots & \vdots \\ 1 & 1 & \dots & 1-x & 1 & \dots & 1 \end{pmatrix}, \quad \begin{cases} x \text{ is real} \\ p+q=n. \end{cases} \quad (3)$$

The rows of the matrix $\mathbf{S}$ are the *generalized* memorized patterns.

The meaningful interpretation of the problem:

the network had to be learned by p -time showing of the standard $\vec{\varepsilon}(n)$,

$$\vec{\varepsilon}(n) = (1, 1, \dots, 1) \in \mathbb{R}^n,$$

but an error crept into the learning process and the network was learned by p distorted copies $\vec{s}^{(l)}$ of the standard:

$$\vec{s}^{(l)} = (1, \dots, 1, \underbrace{1-x}_l, 1, \dots, 1), \quad l = 1, 2, \dots, p. \quad (4)$$

The real number x is called the distortion parameter.

The problem under investigation, Eqs.(1) -(3), is very close to the problem of generalization in the case of one embedded pattern^{2,3}.

Main Results⁴

1°. The local maxima of the functional $F(\vec{\sigma})$ necessarily have the form⁵

$$\vec{\sigma}^* = (\underbrace{\sigma_1, \sigma_2, \dots, \sigma_p}_{\vec{\sigma}'}, 1, \dots, 1), \quad (5)$$

and

$$F(\vec{\sigma}^*) \propto x^2 - 2x(q + p \cos w) \cos w + (q + p \cos w)^2, \quad (6)$$

where w is the angle between vectors $\vec{\sigma}'$ and $\vec{\varepsilon}(p) = (1, 1, \dots, 1) \in \mathbb{R}^p$:

$$\cos w = \frac{\sum_{i=1}^p \sigma_i}{p} = \frac{(\vec{\sigma}', \vec{\varepsilon}(p))}{\|\vec{\sigma}'\| \cdot \|\vec{\varepsilon}(p)\|}. \quad (7)$$

Then, the vectors $\vec{\sigma}^*$ (5) with the p -dimensional parts $\vec{\sigma}'$ equidistant from $\vec{\varepsilon}(p)$, *provide the same value* of the functional F .

2°. Evidently,

$$\cos w \equiv \cos w_k = 1 - \frac{2k}{p}, \quad k = 0, 1, \dots, p,$$

and the vectors $\vec{\sigma}^*$ (5) are grouped into $p + 1$ classes Σ_k on which the functional $F(\vec{\sigma}^*)$ is constant:

$$\boxed{\Sigma_k = \{\vec{\sigma}^* \mid \text{exactly } k \text{ coordinates of } \vec{\sigma}^* \text{ are equal to } "-1"\}.}$$

The number of the vectors $\vec{\sigma}^*$ in the class Σ_k is equal $\binom{p}{k}$.

3°. To find the ground state dependence on x , it is necessary to analyze the family of the straight lines

$$L_k(x) = (q + p \cos w_k)^2 - 2x(q + p \cos w_k) \cos w_k, \quad k = 0, 1, \dots, p. \quad (8)$$

In the region where $L_k(x)$ majorizes all the other straight lines, the ground state belongs to the class Σ_k and it is $\binom{p}{k}$ times degenerated.

4° Theorem.

When x increases from $-\infty$ to ∞ , the ground state in consecutive order belongs to the classes

$$\Sigma_0, \Sigma_1, \dots, \Sigma_{k_{max}}.$$

The k th rebuilding of the ground state ($\Sigma_{k-1} \rightarrow \Sigma_k$) occurs at the point x_k of the intersection of the straight lines $L_{k-1}(x)$ and $L_k(x)$:


$$x_k = p \frac{n - (2k - 1)}{n + p - 2(2k - 1)}, \quad k = 1, 2, \dots, k_{max}, \quad (9)$$

where

$$k_{max} = \min \left(p, \left\lceil \frac{n + p + 2}{4} \right\rceil \right).$$

The functional has no other local maxima.

The Theorem relates the quality of the learning of the network with the value of the distortion x during the learning stage and with the length p of the learning sequence. It is reasonable, that the error of the network increases with the increase of the distortion x : when $x \in (x_k, x_{k+1})$ the class Σ_k ("the truth" understood by the network) differs from the standard $\vec{\epsilon}(n)$ by k coordinates (others interpretations see below).



In the Fig.1 is given the typical behavior of the straight lines $L_k(x)$. The rebuildings of the ground state occurs at the points x_k of the intersection of the straight lines L_{k-1} and L_k . Inside the interval (x_k, x_{k+1}) the ground state belongs to the class Σ_k . When x increases:

- a). all the rebuildings of the ground state occur: $k_{max} = p$;
- b). only $k_{max} = \left\lfloor \frac{n+p+2}{4} \right\rfloor < p$ rebuildings of the ground state occur.

Generalizations of Basic Model⁴

1°. When the standard $\vec{\varepsilon}(n)$ is changed by an *arbitrary* configuration vector $\vec{\alpha} = (\alpha_1, \alpha_2, \dots, \alpha_n)$, $\alpha_i = \{\pm 1\}$, the Theorem remains valid, but the vectors $\vec{\sigma}^*$ (5) have the form

$$\vec{\sigma}^* = (\alpha_1 \sigma_1, \alpha_2 \sigma_2, \dots, \alpha_p \sigma_p, \alpha_{p+1}, \dots, \alpha_n).$$

2°. When we rotate the memorized patterns (4) as a whole, all their first p coordinates are distorted.

Suppose the rotation matrix $\mathbf{U} = (u_{ij})$ transforms the first p coordinates

of n -dimensional vectors only:

$$\begin{cases} \vec{u} = \mathbf{U} \cdot \vec{\varepsilon}(p) = (u_1, u_2, \dots, u_p), \\ u_l = \sum_{i=1}^p u_{li}, \quad l = 1, 2, \dots, p; \quad \|\vec{u}\|^2 = p. \end{cases} \quad (10)$$

Then the memorized patterns take the form:

$$\vec{s}^{(l)} = (u_1 - xu_{1l}, u_2 - xu_{2l}, \dots, u_p - xu_{pl}, 1, \dots, 1), \quad l = 1, 2, \dots, p.$$

It is easy to see, that if the standard $\vec{\varepsilon}(n)$ does not change after the rotation ($u_l \equiv 1$) all the results of the "Basic Model" remain unchanged.

More interesting is the case when the standard $\vec{\varepsilon}(n)$ shifts after the rotation ($u_l \neq 1$). Again the only important configuration vectors are $\vec{\sigma}^*$ (5) and the functional $F(\vec{\sigma}^*)$ is given by the same expression (6). But now w is the angle between vectors $\vec{\sigma}'$ and $\vec{u}$:

$$\cos w = \frac{\sum_{i=1}^p \sigma_i \cdot u_i}{p} = \frac{(\vec{\sigma}', \vec{u})}{\|\vec{\sigma}'\| \cdot \|\vec{u}\|}. \quad (11)$$

The vectors $\vec{\sigma}^*$ are grouped in the classes $\Sigma_k^{(U)}$ on which $F(\vec{\sigma}^*)$ is constant:

$$\boxed{\Sigma_k^{(U)} = \{\vec{\sigma}^* \mid \text{with } p\text{-dimensional parts } \vec{\sigma}' \text{ equidistant from } \vec{u}.\}}$$

The number of the different classes $\Sigma_k^{(U)}$ is given by the number t of *different* values of $\cos w$ (11):

$$\cos w_0 > \cos w_1 > \dots > \cos w_t; \quad \cos w_k = -\cos w_{t-k}, \quad \forall k \leq t.$$

Then we have the following generalization of the Theorem:

when x increases from $-\infty$ to ∞ , the ground state in consecutive order belongs to the classes

$$\Sigma_0^{(U)}, \Sigma_1^{(U)}, \dots, \Sigma_{k_{max}}^{(U)}.$$

The k th rebuilding of the ground state ($\Sigma_{k-1}^{(U)} \rightarrow \Sigma_k^{(U)}$) occurs at the point x_k of the intersection of the straight lines $L_{k-1}(x)$ and $L_k(x)$:

$$x_k = \frac{p}{2} \left[1 + \frac{q}{q + p(\cos w_{k-1} + \cos w_k)} \right], \quad k = 1, 2, \dots, k_{max}. \quad (12)$$

If $x_1 > \frac{3}{4}p$, all the rebuildings take place ($k_{max} = t$). If $x_1 < \frac{3}{4}p$, the rebuildings stop when the denominator in Eq.(12) becomes negative.

Note. The compositions of the classes $\Sigma_k^{(U)}$ are determined by the values of $\{u_l\}_1^p$ only. But the choice of $\{u_l\}$ is completely in our hands! Then we can create the Hopfield type network with a preassigned set of the fixed points.

3°. The memorized patterns can be obtained from $\vec{\varepsilon}(n)$ by *the identical synchronous* distortions of its m coordinates. Suppose $n = p \times m + q$ and the matrix **S** is

$$\mathbf{S} = \begin{pmatrix} 1-x & \dots & 1-x & 1 & \dots & 1 & \dots & 1 & \dots & 1 & 1 & \dots & 1 \\ 1 & \dots & 1 & 1-x & \dots & 1-x & \dots & 1 & \dots & 1 & 1 & \dots & 1 \\ \vdots & \dots & \vdots & \vdots & \dots & \vdots & \dots & \vdots & \dots & \vdots & \vdots & \dots & \vdots \\ 1 & \dots & 1 & 1 & \dots & 1 & \dots & 1-x & \dots & 1-x & 1 & \dots & 1 \end{pmatrix}.$$

The "suspicious" configuration vectors are the piecewise constant vectors

$$\vec{\sigma}^* = (\underbrace{\sigma_1, \dots, \sigma_1}_m, \underbrace{\sigma_2, \dots, \sigma_2}_m, \dots, \underbrace{\sigma_p, \dots, \sigma_p}_m, \underbrace{1, \dots, 1}_q); \quad (13)$$

the values of the functional are

$$F(\vec{\sigma}^*) \propto x^2 - 2x \cos w \left(\frac{q}{m} + p \cos w \right) + \left(\frac{q}{m} + p \cos w \right)^2,$$

where, as in Eq.(7), w is the angle between $\vec{\sigma}' = (\sigma_1, \sigma_2, \dots, \sigma_p)$ and $\vec{\varepsilon}(p)$.

Again the vectors (13) are grouped into classes $\Sigma_k^{(m)}$, whose structure is similar to the structure of the classes Σ_k . Then we have the generalization of the Theorem: *the value of the parameter x , which corresponds to the k th rebuilding of the ground state $(\Sigma_{k-1}^{(m)} \rightarrow \Sigma_k^{(m)})$, is*

$$x_k = p \frac{\frac{n}{m} - (2k-1)}{\frac{n}{m} + p - 2(2k-1)},$$

and

$$k_{max} = \min \left(p, \left\lceil \frac{\frac{n}{m} + p + 2}{4} \right\rceil \right).$$

Basic Model: Sequences and Interpretations⁴

$$\boxed{1^\circ. \quad x_1 = p \frac{n-1}{n+p-2} \geq \frac{p}{2}.}$$

In fact x_1 is the boundary of the distortions up to which the network reproduces the standard from its distorted copies correctly. The boundary depends on the length p of the learning sequence. Of course, the network is learned correctly, if the value of the distortions does not exceed $\frac{p}{2}$.

$$\boxed{2^\circ. \quad x_1 \text{ is monotonically increasing function of } p \text{ and } n.}$$

Let n and x be fixed. Merely due to an increase of p the boundary x_1 can be forced to exceed x (if x is not too large). As a result x turns out to be on the left of a new position of x_1 , i.e. in the region where *the only* fixed point is the standard $\vec{\varepsilon}(n)$. In other words, only by an increase of the length p of the learning sequence we can force the network to understand correctly "the truth" it is tried to be learned. It is in agreement with the practical experience: the greater the length of the learning sequence, the better the signal can be read through noise.

Let p and x be fixed. As above, merely due to an increase of the number n the value of x_1 can be forced to exceed x . This result is reasonable too: if p is fixed, the greater is n , the smaller is the relative weight $\frac{p}{n}$ of the distorted coordinates. Naturally, the less is the relative distortion, the better must be the result of the learning.

3°. $x_{\frac{p+1}{2}} = p$. (Without the loss of generality we assume that p is odd.)

Here we change the notation for the standard $\vec{\varepsilon}(n)$ and introduce another standard $\vec{\varepsilon}^{(-)}$:

$$\begin{aligned}\vec{\varepsilon}^{(+)} &= (1, 1, \dots, 1, 1, \dots, 1) = \vec{\varepsilon}(n) \\ \vec{\varepsilon}^{(-)} &= (-1, -1, \dots, -1, 1, \dots, 1).\end{aligned}$$

In their not coincident parts the standards $\vec{\varepsilon}^{(+)}$ and $\vec{\varepsilon}^{(-)}$ are opposed with each other, i.e. they are two opposite "statements". Any of the network fixed points $\vec{\sigma}^*$ (5) is an intermediate statement between $\vec{\varepsilon}^{(+)}$ and $\vec{\varepsilon}^{(-)}$, which is drawn towards either one edge of the scale, or the other. And the network "feels" this.

Indeed, when the distortion x is not very large ($x < p$), the number k of the ground state does not exceed $\frac{p}{2}$, and the ground state more resembles $\vec{\varepsilon}^{(+)}$ than $\vec{\varepsilon}^{(-)}$. In other words, the memorized patterns are interpreted by the network as the distorted copies of the standard $\vec{\varepsilon}^{(+)}$. But if during the learning stage the distortion exceeds p ($x > p$), the number of ground state exceeds $\frac{p}{2}$ and the ground state resembles $\vec{\varepsilon}^{(-)}$. Now the network interprets the memorized patterns as the distorted copies of another standard $\vec{\varepsilon}^{(-)}$.

This is in agreement with the practical experience: we interpret deviations in the image of a standard as permissible *only up to some boundary*. If only this boundary is exceeded, the patterns are interpreted as the distortions of quite different standard. For the network of the considered type this boundary is p .

One extra argument to support this interpretation: from Eq.(9) it is easy to see that when $p = \text{const}$ and $n \rightarrow \infty$ all x_k stick to one point

$$x_k \equiv p, \quad k = 1, 2, 3, \dots, p;$$

then for $x < p$ the ground state belongs to the class $\Sigma_0 = \vec{\varepsilon}^{(+)}$, whereas for $x > p$ the ground state belongs to the class $\Sigma_p = \vec{\varepsilon}^{(-)}$.

$$4^\circ. \quad k_{max} = \begin{cases} p, & \text{when } \frac{p-1}{n-1} < \frac{1}{3} \\ \left[\frac{n+p+2}{4} \right], & \text{when } \frac{p-1}{n-1} > \frac{1}{3} \end{cases}$$

So, p is the boundary for the permissible distortions x . The question is, what do the memorized patterns with large distortions x mean? We treat the increase of x above p as the more and more negation of the standard $\bar{\varepsilon}^{(+)}$. As if the network is learned by the memorized patterns, which deny the standard $\bar{\varepsilon}^{(+)}$. In other words, the network is *relearned by presentation of negative examples*.

There is big and clear to everybody difference between the relearning with the help of negative examples and *the learning of the opposite truth*. The relearning is characterized by some specific difficulties: (1) the better the incorrect truth has been understood, the more difficult (and sometimes even impossible) to correct it; (2) it is comparatively easy to correct the result slightly, but it is much more difficult to revise it in the main, *etc.* We think, that the dependence of k_{max} on p is the reflection of just these problems.

When the number p of the parameters which have to be corrected is not very great ($\frac{p-1}{n-1} < \frac{1}{3}$), the network can be relearned by simple presentation of negative examples. In this case $k_{max} = p$ and, when "the denial" of the standard $\bar{\varepsilon}^{(+)}$ is rather strong ($x > x_p$), as "a new" truth the network understands the opposite standard $\bar{\varepsilon}^{(-)}$. But if the number of the corrected parameters is great ($\frac{p-1}{n-1} > \frac{1}{3}$), to relearn the network it is not sufficient to present the negative examples. In this case $\left[\frac{p+1}{2} \right] < k_{max} < p$ and whatever large x is ($x_{k_{max}} < x < \infty$), as a new truth the network understands not the opposite standard $\bar{\varepsilon}^{(-)}$, but one of the statements intermediate between $\bar{\varepsilon}^{(+)}$ and $\bar{\varepsilon}^{(-)}$. Though the understood truth is drawn towards $\bar{\varepsilon}^{(-)}$, since $k_{max} > \frac{p}{2}$.

Of course, our interpretation is open for discussion. But it seems that in real life there are a lot of examples, which confirm our conception.

New Results (in preparation)

The generalization of the Basic Model to the cases:


1°. The functional $F(\vec{\sigma})$ in the problem (1)-(3) has the form

$$F(\vec{\sigma}) = \sum_{i,j=1}^n J_{ij} \sigma_i \sigma_j + h \sum_{i=1}^n \sigma_i.$$

In physics such a linear term describes the magnetic field.

In the Fig.2 the straight lines $h_k(x)$ divide the plane (x, h) into the regions where the ground state belongs to the different classes Σ_k ,

$$h_{k-1}(x) = 2p(n - 2k + 1) \left(\frac{x}{x_k} - 1 \right), \quad k = 1, 2, \dots$$



2°. The distortions x_l are different for all the memorized patterns (4):

$$\vec{s}^{(l)} = (1, \dots, 1, \underbrace{1 - x_l}_l, 1, \dots, 1), \quad l = 1, 2, \dots, p.$$

Suppose

$$x_1 \geq x_2 \geq \dots, \geq x_p > 0.$$

It can be shown that, firstly, only one of the configuration vectors

$$\vec{\sigma}^*(k) = (\underbrace{-1, -1, \dots, -1}_k, 1, \dots, 1), \quad k = 0, 1, \dots, k_{max},$$

can be the ground state. Here

$$k_{max} \begin{cases} = p, & \text{when } p < \frac{n}{2} \\ \leq \left[\frac{n}{2} \right], & \text{when } p > \frac{n}{2}. \end{cases}$$

And secondly, **in order to the vector $\vec{\sigma}^*(k)$ be a ground state, the fulfilment of the inequalities**

$$x_{k+1} - \frac{\sum_{i=1}^k x_i - \sum_{j=k+2}^p x_j}{n - 2k - 1} \leq p \leq x_k - \frac{\sum_{i=1}^{k-1} x_i - \sum_{j=k+1}^p x_j}{n - 2k + 1}$$

is necessary and sufficient conditions.

When p is fixed and $n \rightarrow \infty$ these inequalities are much more simpler:

$$x_{k+1} \leq p \leq x_k.$$

Note, that for sufficiently large n and $p > x_1$, the standard $\vec{\varepsilon}(n) = \vec{\sigma}^*(0)$ is the ground state. It seems, that for the Hebb connection matrix the last result clarifies the meaning of the well-known Latin saying "*Repetitio est mater studiorum*" – showing the same pattern many times (inevitably each time with a distortion), we seek to make the number of the presentations p greater than the maximal distortion.

¹ Litinskii L.B. 1997 Energy functional and fixed points of neural network. cond-mat/9706280. Also in: M.Marinaro and R.Tagliaferri (Eds.) 1998 "Neural Nets. WIRN VIETRI-97. Proceedings..." (Berlin: Springer-Verlag) p.338.

² Fontanari J.F. 1990 J. Phys. France v.51 p.2421.

³ Krebs P.R., Theuman W.K. 1993 J. Phys. A26 p.3983.

⁴ Litinskii L.B. 1999 Theor. and Math. Phys. v.118 p.133. cond-mat/9906197.

⁵ Litinskii L.B. 1994 Theor. and Math. Phys. v.101 p.1492.